

## SOME EMBEDDING THEOREMS FOR MODAL LOGIC

DAVID MAKINSON

We\* shall prove some embedding theorems for modal logic, that is, theorems to the effect that every consistent modal logic satisfying certain general conditions is a sublogic of certain particular logics. Our results are related to those of McKinsey [1] and Tarski [2].

We begin with terminology. *Formulae* are understood to be built from a denumerable list of elementary letters by means of the operators  $\neg$ ,  $\wedge$ ,  $\square$ , with other operators introduced as usual. By a *modal logic* we mean any set  $S$  of formulae that contains all the tautologies in  $\neg$  and  $\wedge$  and is closed under the operations of substitution (of arbitrary formulae for elementary letters) and detachment ( $\alpha, \alpha \supset \beta / \beta$ ). We say that a set  $S$  of formulae is closed under *congruence* if whenever  $(\alpha \equiv \beta) \in S$  then  $(\square \alpha \equiv \square \beta) \in S$ ; closed under *monotony* if whenever  $(\alpha \supset \beta) \in S$  then  $(\square \alpha \supset \square \beta) \in S$ ; closed under *antitony* if whenever  $(\alpha \supset \beta) \in S$  then  $(\square \beta \supset \square \alpha) \in S$ .

By a *modal algebra* we mean a structure  $\mathfrak{A} = \langle A, -, \cap, * \rangle$  where  $\langle A, -, \cap \rangle$  is a Boolean algebra and  $*$  is a unary operation over  $A$ . A modal algebra is said to be *monotonic* if for all  $x, y \in A$ ,  $x \leq y$  implies  $*x \leq *y$ , and is said to be *antitonic* if for all  $x, y \in A$ ,  $x \leq y$  implies  $*y \leq *x$ . Among the modal algebras there are clearly just four that can be obtained by adding a unary operation to the two-element Boolean algebra: we shall call these the *unit* algebra ( $*1 = 1, *0 = 1$ ), the *identity* algebra ( $*1 = 1, *0 = 0$ ), the *complement* algebra ( $*1 = 0, *0 = 1$ ), and the *zero* algebra ( $*1 = 0, *0 = 0$ ). Each of these four algebras determines a corresponding set of formulae, consisting of just those formulae that are valid in the algebra, that is, just those formulae  $\alpha$  such that for every homomorphism  $h$  from formulae into that algebra,  $h(\alpha) = 1$ . It is easy to verify that each of these four sets of formulae is a modal logic in the sense defined, is closed either under monotony or under antitony, and can be axiomatized in a trivial way: we refer to these four sets of formulae as the *unit*, *identity*, *complement*, and *zero* modal logics respectively.

---

\*Work for this paper was carried out while the author was on contract with the Organization of American States, Department of Scientific Affairs.

**Theorem 1.** *Let  $S$  be any consistent modal logic that is closed under congruence and contains the theses  $\Box(p \vee \neg p)$  and  $\neg\Box(p \wedge \neg p)$ . Then  $S$  is a sublogic of the identity logic.*

*Proof:* We use an algebraic argument. Define a relation over formulae by putting  $\alpha \simeq \beta \pmod{S}$  if  $(\alpha \equiv \beta) \in S$ . It can be verified that since  $S$  is a modal logic in the sense defined, the relation is an equivalence and is congruential with respect to the operators  $\neg$  and  $\wedge$ . Further, since  $S$  is closed under congruence, the relation is clearly congruential with respect to the modal operator  $\Box$ . Hence we can form a Lindenbaum algebra  $|S|$  of  $S$  as the quotient structure determined by the relation. It can be verified that  $|S|$  is a modal algebra  $\langle A, -, \cap, * \rangle$  which, since  $S$  is consistent, has at least two elements. Also, for every formula  $\alpha$ ,  $\alpha \in S$  if and only if  $\alpha$  is valid in  $|S|$ —that is, if and only if  $h(\alpha) = 1$  for every homomorphism from formulae into  $|S|$ .

Moreover, since  $\Box(p \vee \neg p) \in S$  and  $\neg\Box(p \wedge \neg p) \in S$  we have  $*1 = 1$  and  $*0 = 0$ . Thus the set  $\{1, 0\}$  consisting of the unit and zero elements of  $|S|$  is closed under all three operations  $\neg$ ,  $\wedge$ ,  $\Box$ , and so forms a subalgebra of  $|S|$  which clearly coincides with the identity algebra. Now on universal algebraic grounds every formula that is valid in  $|S|$  is valid in all of its subalgebras. So since  $|S|$  is characteristic for  $S$ , we have that  $S$  is a sublogic of the identity logic.

**Theorem 2.** *Let  $S$  be any consistent modal logic that is closed under monotony. Then  $S$  is a sublogic of the identity logic, or the zero logic, or the unit logic.*

*Proof:* Clearly any modal logic that is closed under monotony is closed under congruence, and we can form a characteristic Lindenbaum algebra in the same way as in the proof of theorem 1. At this point the argument divides into three cases.

Case 1. Suppose that  $\Box(p \vee \neg p) \in S$  and  $\neg\Box(p \wedge \neg p) \in S$ . Then the conditions of theorem 1 are satisfied and so  $S$  is a sublogic of the identity logic.

Case 2. Suppose that  $\Box(p \vee \neg p) \notin S$ . Then  $*1 \neq 1$  and so by standard results on Boolean algebras there is an ultrafilter  $X$  of  $|S|$  with  $*1 \in X$ . Now it can be verified that since  $S$  is closed under monotony,  $|S|$  is monotonic. Thus for all  $x \in |S|$  we have  $x \leq 1$  and so  $*x \leq *1$  and so  $*1 \leq *x$  and so  $*x \in X$  and so  $*x \notin X$ . We define a function  $h$  from  $|S|$  into the zero algebra as follows: if  $x \in X$  put  $h(x) = 1$ , and if  $x \notin X$  put  $h(x) = 0$ . Since  $X$  is an ultrafilter,  $h$  is homomorphic with respect to the Boolean operations. Further, for each  $x \in |S|$  we have  $h(*x) = 0 = *h(x)$  and so  $h$  is a homomorphism from  $|S|$  into, and indeed clearly onto, the zero algebra. Now on universal algebraic grounds every formula that is valid in  $|S|$  is valid in all of its homomorphic images, and so since  $|S|$  is characteristic for  $S$ , we have that  $S$  is a sublogic of the zero logic.

Case 3. Suppose that  $\neg\Box(p \wedge \neg p) \notin S$ . Then  $*0 \neq 0$ , and so by standard results on Boolean algebras there is an ultrafilter  $X$  of  $|S|$  with  $*0 \in X$ . We can use an argument similar to that of the second case to show that  $S$  is a sublogic of the unit logic.

**Theorem 3.** *Let  $S$  be any consistent modal logic that is closed under antitony. Then  $S$  is a sublogic of the complement logic, or the unit logic, or the zero logic.*

*Proof:* We can use the same kind of argument as for theorem 2.

#### REFERENCES

- [1] McKinsey, J. C. C., "On the number of complete extensions of the Lewis systems of sentential calculus," *The Journal of Symbolic Logic*, vol. 9 (1944), pp. 42-45.
- [2] Tarski, A., "On extensions of incomplete systems of the sentential calculus," in *Logic, Semantics, Metamathematics*, transl. by J. Woodger, Oxford: Clarendon Press (1956).

*American University of Beirut  
Beirut, Lebanon*